

Helping Asthmatic Children Through Bonding Therapy

**Antonio Madrid, PhD, Dale Pennington, PhD,
Gary Brown, PhD & Maureen Wolfe, MA**

Abstract: Disruptions in maternal-infant bonding are shown to be the mediating variable between maternal distress and the subsequent expression of childhood asthma. When bonding is repaired, it seems that children's asthmatic symptoms diminish or remit. This study evaluated 16 asthmatic children before and after their mothers were treated with bonding therapy. Fourteen improved on 11 measures, including reduction in STEP classification system and medication use. Thirteen children were able to stop all medications. The links between bonding disruptions, stress, and airway inflammation are discussed.

Keywords: Bonding Therapy, Childhood Asthma, Bonding Disruption, Asthma Treatment

Introduction

Asthma affects more children's lives than any other chronic illness (Lara et al., 2002). In 2008, just over 10 million children within the U.S were reported as ever having had a diagnosis of asthma (Akinbami, Moorman, Garbe, & Sondik, 2009). It is the third-ranking cause of hospital admissions in children under 15 (DeFrances, Cullen, & Kozak, 2007). The financial cost in America in 2007 was \$56 billion in health costs, lost school and work days, and early deaths (Center for Disease Control and Prevention, 2011). Asthma is responsible for 14 million missed school days annually (Martin, 2004)—more than any other chronic illness.

Since childhood asthma has reached epidemic proportions and has major public health and financial impacts, researchers have been attempting to identify risk factors for the development of this disease. For several years a significant focus of attention has revolved around the connection between maternal distress and the subsequent

The authors wish to thank the Association of Pre and Perinatal Psychology and Health, The Mental Insight Foundation, and the Huizenga Family for their generous contributions to this study. Correspondence can be sent to Antonio Madrid, PhD, POB 519, Monte Rio, CA 95462, madrid@sonic.net.

development of asthma children.

Mrazek, Klinnert, Mrazek, and Macey (1991), at the National Jewish Center for Immunology and Respiratory Medicine in Denver, studied 150 children whose mothers were asthmatic. They found a link between early problems in coping/parenting and the subsequent expression of asthma. Klinnert et al. (2001), surprised by this finding, noted that this link between parental stress in care-giving and the subsequent development of asthma was the first documented report of such a connection.

Maternal stress is known to be caused by many factors: Cesarean section deliveries, maternal health issues, psychological problems, maternal despondency, to list a few. Cesarean section deliveries and emergency cesarean sections have been linked to childhood asthma. A Finnish study of 60,000 births found that mothers who delivered by cesarean sections were 50% more likely to have a child who later developed asthma (Kero et al., 2002). Emergency cesarean section deliveries (even more stressful) predictably raised the asthma rate up to 60%. This finding was replicated by Roduit et al. (2009) who studied 2,917 children.

Annesi-Maesano, Moreau, and Strachan (2001), using a British cohort of 2,583 mothers, investigated whether in utero and perinatal influences contribute to the development and severity of asthma in childhood. Childhood asthma was more frequently reported by mothers when there had been health complications during pregnancy, labor, or delivery, or when the child was ill during the first week of life. She concluded that there is evidence that in utero and perinatal factors may increase the risk of developing asthma. Similarly, a Norwegian study of over 1.5 million mothers and 5,938 asthmatic children found that many types of pregnancy complications represented a risk factor for the development of asthma in the offspring (Nafstad, Samuelsen, Irgens, & Bjerkedal, 2003).

Kozyrskyj et al. (2008) studied healthcare records of 13,907 children and their mothers from Manitoba databases. Healthcare or prescription medication for depression or anxiety was used to define maternal distress, and asthma status was determined from the children's asthma prescription records. They found that risk for childhood asthma was increased among children who were exposed to continued maternal distress from birth until age 7 years. Similarly, a Puerto Rican study concluded that maternal depressive symptoms were associated with an increased risk of asthma hospitalizations at age 1 year (Lange et al., 2011).

Mother-child interactions have been shown to predict the

development of asthma in the child by school age. Mantymaa et al. (2003) showed that psychological stress is associated with physical illnesses like asthma or infection. As mentioned above, Klinnert et al. (2001) investigated many factors associated with childhood asthma and found that global assessment of parenting problems was a predictor of the development of asthma by age 3.

In that same vein, Wright, Cohen, Carey, Weiss, and Gold (2002) found that greater levels of caregiver-perceived stress at 2 to 3 months was associated with increased risk of subsequent repeated wheeze among children during the first 14 months of life. Further, they found that prenatal stress was associated with altered innate and adaptive immune responses, concluding that stress-induced perinatal immunomodulation may impact the expression of allergic disease in these children (Wright et al., 2010).

Cassibba, van Ijzendoorn, Bruno, and Coppola (2004) found that children affected by asthmatic bronchitis were less securely attached than healthy comparisons. These children showed less harmonious and comfort seeking behaviors than healthy children, indicating insecure attachment.

Three studies investigated the relationship between pediatric asthma and the difficulties that a mother experiences in bonding with her baby. Using the "maternal-infant bonding" paradigm of Klaus and Kennell (1976), these studies looked at the incidence of bonding problems within pediatric asthma populations as compared to well-baby groups. Klaus and Kennell had previously demonstrated that a mother's love for her infant will likely be disrupted if the mother and infant were separated at birth or if the mother was experiencing some great stress or trauma in her life during the pregnancy or the perinatal period.

In the first of these studies Feinberg (1988) showed that bonding disruptions occurred three times more frequently in mothers of asthmatics than in mothers of well-babies (84% vs 24%). Schwartz (1988) found almost identical numbers: 86% vs 29%.

Pennington (1991), in the third study, found that four "non-bonding events" were most predictive of asthma: delay in holding the baby, family death in first year, emotional problems during pregnancy, and maternal emotional problems in the first year. He concluded that bonding disruptions appear to be the mediating variable that links pediatric asthma with the various maternal factors and stressors that have been identified by numerous researchers.

If this is the case—that bonding disruptions are the link between all of these maternal stressors and pediatric asthma—the next

question logically surfaces: "What happens if the mother becomes bonded to her asthmatic child?"

From our clinical experience over 15 years, we had observed that a child's asthma seemed to improve once a disrupted bond between a mother and child was repaired. We subjected this observation and hypothesis to two pilot clinical trials. In these studies, each of the mothers was treated with a therapy that focused entirely on processing the traumas surrounding the birth of her child and then creating an imagined idealized birth. We hypothesized that this would repair the disrupted maternal-infant bond and would have an impact on the child's asthma.

The first study involved six mother-child pairs (Madrid, Ames, Skolek, & Brown, 2000). Five of the six children, including two infants, experienced complete or nearly complete remission from asthma symptoms as measured across 18 variables. In the second study (Madrid, Ames, Horner, Brown, & Navarrette, 2004), asthma symptoms improved in 12 of 15 children. Eight of the 10 children who were taking medication no longer needed to continue them.

The present study continues the investigation of the hypothesis that children's breathing problems can be improved by working with the mothers to repair the broken maternal-infant bond. This study focuses on the use of measures that are more objective.

Method

Sample

Eighteen asthmatic children and their mothers were identified for this study, from responses to newspaper and magazine articles, posted flyers and mailings, and referrals from other therapists. There were three criteria for inclusion in the sample. (1) The child's asthma needed to be diagnosed by a physician. (2) The mother had to be psychologically stable, having a GAF score of 70 or better. GAF (Global Assessment of Functioning) is a scale from the *Diagnostic and Statistical Manual*, used by mental health workers to assess psychological disorders (American Psychiatric Association, 2000). (3) The mother needed to have a pre or perinatal history that met the criteria for non-bonding.

The diagnosis of asthma is always problematic because asthma is not a homogeneous condition and is often over-diagnosed (Martinez, 2002). While asthma is a disease consisting of airway constriction and inflammation, it is not unusual for viral wheezing, atopic reactivity,

and bronchial inflammation to be diagnosed as asthma (Kozyrskyj, Mustard, & Becker, 2003). Rather than entering into a debate about the accuracy of the diagnosis of our subjects, we simply decided to use a physician's diagnosis as the criterion for admission into the cohort.

Of the original 18 subjects, one was eliminated because the mother did not meet the GAF standard, and one was eliminated because the therapy deviated from the prescribed standard. All of the mothers in the initial sample had prenatal or perinatal histories consistent with problems in bonding, i.e., there was evidence of either separation at birth or maternal trauma during pregnancy or soon afterwards.

The remaining 16 children ranged in age from 9 months to 19 years. All were under the care of a physician and all were taking medication.

Procedures

Following the mother's recruitment into the study, an initial interview was conducted either in person or by telephone. *The Maternal-Infant Bonding Survey* (MIBS) (Brown, Pennington, & Madrid, 1999) was administered to determine if she met the criterion of a mother with disrupted bonding. The *MIBS* (see Appendix A), a survey of events which have been linked to bonding disruptions, was developed from the work of Klaus and Kennell (1976).

The mother was then given a questionnaire (see Appendix B) which asked:

- the age of her child when asthma was first diagnosed and the circumstances under which it was diagnosed;
- the current age of the child;
- the way the child's asthma presented itself, including triggers;
- her assessment of the severity of her child's asthma; and
- specific measures of her child's condition (described in the Measures section below).

She was then administered the Beck Depression Inventory.

After being given a description of the study, including the treatment proposed, the mothers began treatment, described in the *Treatment* section below.

Measures

Eleven variables were used to measure each child's condition before and six months after treatment.

1. Asthma Monitor—a five question instrument which assesses current asthma severity. Each answer was scored from 0 to 4.

The five scores were then added for the child's total score. Higher numbers indicate greater severity. Total scores range from 0 to 20. (See Appendix B.)

2. STEP—used by the National Institutes of Health (U.S. Department of Health and Human Services, 2003) to determine asthma severity and to design treatment protocol (see Appendix B). Two questions are asked about “daytime symptoms” and “night time symptoms,” which place the child into one of five Steps or categories:

No Asthma Symptoms (0)

Mild Intermittent (1)

Day Time Symptoms: equal to or less than 2 days/week
 Night Time Symptoms: equal to or less than 2
 night/week

Mild Persistent (2)

Day Time symptoms: greater than twice/week but less
 than once daily
 Night time Symptoms: greater than 2 nights/month

Moderate Persistent (3)

Day Time Symptoms: daily
 Night Time Symptoms: more than 1 night/week

Severe Persistent (4)

Day Time Symptoms: continual
 Night time Symptoms: frequent

3. Doctor Visits: number of unscheduled visits during last 6 months.
4. Emergency Room Visits - number of ER visits during last 6 months
5. Hospital stays - number of admissions and total days during prior 6 months.
6. Oral steroids – the number of courses and total number of days during the prior 6 months.
7. Days housebound – number of days that child kept from school or stayed in bed or could not go outside and play during prior 6 months.
8. Asthma symptoms with exercise—whether the child wheezes when he is active, rated on a four point scale from Never (0) to Always (3).

9. Overall Health – mother’s perception of the child’s general health. Scored: 0 to 4.
10. Energy Level – mother’s perception of the child’s vitality. Scored 0 to 4
11. Medications – list of the medications that the child was currently taking.

All mothers were asked to take the Beck Depression Inventory-II (BDI) immediately before treatment and 6 months following treatment. The BDI is a 21 item, multiple choice, self-report inventory, widely used by mental health professionals to assess the severity of a patient’s depression (Beck, Ward, Mendelson, Mock, & Erbaugh, 1961). Each person’s total score can range from 0 to 63, with 0-5 considered a low score, 6-14 a medium score, and 15-and-above a high score.

Treatment

Each mother was seen in therapy, using a protocol designed to repair the disrupted bond that existed at or near the time of her child’s birth. The treatment used a three phase approach.

1. Phase One: With the mother, identify all Non-Bonding Events (NBE) i.e., the events or situations that interfered with bonding. This was accomplished using the Maternal Infant Bonding Survey and by discussing with the mother her pregnancy and the birth of her child.
2. Phase Two: Afford the mother an opportunity with hypnosis to process each NBE until the impact of that event was no longer recorded as painful or disturbing. A detailed description of this procedure can be found in *American Journal of Clinical Hypnosis* (Madrid, 2005/2006). Other interventions besides hypnosis can be used, such as EMDR, art work, Gestalt Therapy, psychodrama (Madrid, Skolek, & Shapiro, 2006).
3. Phase Three: Help the mother to imagine a new birth experience, without the interference of the NBE.

Most mothers were able to satisfactorily complete the treatment in two sessions or less.

Results

Paired-samples t-tests were used to assess improvement in 11 variables after Bonding Therapy. A one-sided P value < 0.05 was considered statistically significant. Means are reported \pm 1 SD.

A total of 16 patients were enrolled into the study. Patients ranged

Table 1. MATERNAL-INFANT BONDING RESULTS

VARIABLE	PRE-TREATMENT	POST-TREATMENT	P VALUE
Asthma Monitor	6.7±4.4	1.8±2.5	< 0.001
STEP	3.4±1.4	0.7±1.1	< 0.001
Doctor visits	1.8±1.8	0.2±0.5	< 0.001
ER visits	0.4±1.0	0.0±0.0	0.055
Days in hospital	0.1±0.3	0.0±0.0	NS*
Oral steroid course	0.8±1.1	0.3±0.9	0.036
Days housebound	6.4±7.0	0.5±1.3	0.003
Asthma symptoms with exercise	1.7±1.5	0.6±1.1	0.018
Health Evaluation	1.1±1.0	0.5±0.7	0.006
Energy Evaluation	0.9±1.1	0.3±0.5	0.020
Mother's Depression Beck Depression Inventory (BDI)	16.2±10.6	3.5±5.5	< 0.001

* NS denotes not significant

in age from 0.75 to 19 years (mean age ranged from 8.0±4.7 years). Age of asthma diagnosis ranged from 0.2 to 9 years (mean age ranged from 4.0±3 years). Fifty-six percent of the patients were male.

The results of Bonding Therapy indicated significant improvement in children's asthma following treatment of the mothers to improve bonding.

An analysis of each of the variables used to measure the effects of treatment herein follows.

Measures

Asthma Monitor, a measure for assessing asthma severity, was collected for 15 patients. There was significant improvement in the pre-treatment 6.7±4.4 versus post-treatment 1.8±2.5 average scores ($P < 0.001$). Every child except one of the two oldest children improved in his Asthma Monitor score.

STEP, a national standard for assessing asthma severity, registered significant improvement in the 15 patients measured, with average pre-treatment scores of 3.4±1.4 compared with the post-treatment average 0.7±1.1 ($P < 0.001$). The average score before treatment was between Moderate Persistent to Severe Persistent; after treatment the average score was between No Asthma to Mild Intermittent. Before treatment, there were no children in the No Asthma class: post treatment there were 9 children in this class. Before treatment there were 8 children with Severe Persistent asthma, and by the end of treatment, there were none in this category.

Except for the two oldest children, every child's STEP classification improved after treatment.

Unscheduled Doctor Visits were reported by 10 of the 16 patients before the study, resulting in 1.8 ± 1.8 mean visits. Only one patient went to the doctor after bonding therapy 0.2 ± 0.5 ($P < 0.001$).

ER visits before treatment was reported by 4 out of 16 patients for an average of 0.4 ± 1.0 visits. No patient went to the ER post-treatment 0.0 ± 0.0 ($P=0.055$).

Hospitalizations were reported by one child who spent 1 day in the hospital pre-treatment 0.1 ± 0.3 . No patients were hospitalized after treatment. The sample size is insufficient to establish significance.

Oral Steroids were used by 7 patients before treatment, varying from 1 to 3 courses for an average of 0.8 ± 1.1 courses. Post-treatment, two patients required (a 2 and 3 course) oral steroid treatment 0.3 ± 0.9 ($P=0.036$).

Days Housebound were experienced by 8 patients pre-treatment (4 ± 7.0 days) and by 2 patients post-treatment (0.5 ± 1.3 days, $P=0.003$).

Asthma symptoms from exercise were reduced from (1.7 ± 1.5 to 0.6 ± 1.1 , $P=0.018$).

Overall Health, as assessed by mothers improved after treatment (1.1 ± 1.0 versus 0.5 ± 0.7 , $P=0.006$).

Energy Level as assessed by mothers also improved after bonding therapy (0.9 ± 1.1 versus 0.3 ± 0.5 , $P=0.020$).

Medication. All 16 patients began the study on a variety of asthma maintenance medications, with some patients taking up to three different prescriptions. Post-treatment, 13 patients no longer required maintenance medication. Two of the three patients who remained on a medication were the older adolescents.

Beck Depression Inventory. In addition to the children's measurements, the mother's depression as scored on the Beck Depression Inventory (BDI) was collected before and after bonding therapy for 13 mothers. Prior to treatment the scores ranged from 3 to 37 (mean= 16.2 ± 10.6). Significant improvement was observed post-treatment with a range of 0 to 16 (mean= 3.5 ± 5.5 , $P < 0.001$).

All of the mothers DBI-II scores improved following treatment, even those of the mothers of the two older children, whose asthma showed no improvement. Two mothers jumped from being severely depressed to either mildly or minimally so. Six mothers changed from being mildly or moderately to minimally depressed. Four mothers remained minimally depressed, although their scores were lower after treatment. Three mothers did not take the BDI-II.

Table 2

Beck Depression Inventory—II
Scores Before and After Treatment

Pt ID	Age of child	Pre	Severity	Post	Severity
301	10	14	Mild	2	Minimal
302	4.5	16	Mild	2	Minimal
303	0.75	37	Severe	7	Minimal
304	9	16	Mild	0	Minimal
306	4.5	8	Minimal	0	Minimal
307	11	16	Mild	2	Minimal
309	6.5				
313	19	3	Minimal	0	Minimal
314	15	3	Minimal	0	Minimal
315	7	16	Mild	0	Minimal
321	9				
323	2.5				
327	9	18	Mild	14	Mild
328	6.5	35	Severe	15	Mild
329	3	6	Minimal	1	Minimal
331	11	23	Moderate	1	Minimal

Discussion

This study, using more stringent measures of asthma symptoms, strengthens our previous findings that bonding a mother to her child seems to improve the child's asthma symptoms. With the exception of the two oldest adolescents, every child in the study showed improvement in all five categories of the Asthma Monitor: getting work done; shortness of breath; awakening at night; use of rescue inhaler; asthma out of control.

Every child in the study (except the two older adolescents) improved in the STEP measure of asthma severity, with the average moving down from "Moderate Persistent or Severe Persistent" to "No Asthma or Mild Intermittent." This improvement occurred without working directly with any of the children themselves but only with the mother.

There were fewer unscheduled doctor visits, fewer trips to the emergency room, and every child had fewer housebound days.

As we had discovered in our previous studies, this treatment does not seem to work for older adolescents. There has been speculation that this may be that bonding improvement has little effect on adolescents who are developmentally in the process of trying to separate from their parents.

We were surprised to find that bonding therapy improved the emotional state of every mother in the study, even mothers of the two older adolescents who did not improve. Some mothers, who were greatly depressed, improved considerably, with no other treatment than repairing the bond. None of the mothers received psychotherapy, but they responded as if they had. It appears that disrupted bonding may be responsible for some amount of depression in mothers, and that this is alleviated when the bond is repaired.

Because of concerns that asthma is often over-diagnosed, it would be useful in future studies to examine the incidence of bonding disruptions as well as the effectiveness of this treatment in all the different presentations of breathing problems in children.

The development of asthma has been linked to several types of maternal distress: psychological stress, health complications, mental health issues, care-giving problems, cesarean section deliveries, and pregnancy complications, among others. When examined through the lens of a maternal-infant bonding, they all fit the paradigm of "non-bonding events" (Pennington, 1991). Each of these can create in the mother an emotional state that is incompatible with bonding and can reduce the probability that bonding will occur (Klaus & Kennell, 1976).

It is our hypothesis that the mediating variable between these maternal distresses and asthma is the disruption in bonding between the mother and child. This hypothesis appears to be confirmed when we notice that the breathing problems improve once bonding is repaired.

How do disruptions in bonding lead to the subsequent expression of asthma and related breathing problems? Pennington (1991) addressed this question when he reviewed the relationship between asthma and maternal issues. Summing up the literature in this area, he offers a hypothesis that a child's lack of connection with his mother "generates fear and its related behaviors, including endocrine and autonomic responses which lead to release of mediators responsible for broncho-constriction" (p. 163).

Mead (2004) in reviewing separation upon the immune system and organ development writes:

Separation in early life is associated with changes in hypothalamic-pituitary-adrenal (HPA) responses to stress, transient and long-term changes in immune competence in non-human primates, and reduced maternal-infant attunement. The impact of maternal-infant separation during the sensitive period may permanently alter affectional ties, and may consequently influence developing organ systems, including the nervous system. (p. 5)

Klennert et al. (2001) state that immune functioning in adults is known to be affected by stress and that this may alter the course of immune-based diseases such as allergies and asthma. So, too, parenting difficulties (i.e., resulting from bonding problems, in our way of thinking) may affect infants the way that stress affects adults and lead to a change in certain aspects of infant's immune system. Klennert hypothesizes that the quality of care-giving affects the emotional and physiologic regulation of the infant and "could alter the hypothalamic-pituitary-adrenal system or immune functioning in the direction of increased allergic response" (p. 7).

Milam et al. (2008) write that the link between maternal stress and childhood stress is well known, and it seems apparent that childhood stress has a bearing on the endocrine system, the autonomic control of airways, and the immune system. Stress leads to the release of hormones that influence the immune system. Stress is linked to the secretion of adrenocorticotrophic hormone and cortisol, which increases the production of inflammatory cytokines. Therefore, it is

possible that stress in a child can alter the functioning of the immune system which then causes inflammatory responses to triggers in the environment.

What greater stress could there be for a baby than to be chronically disconnected from its mother? We think that this stress in the unbonded child can lead, in some children, to inflammatory airway responses; and the present study seems to confirm this link.

References

- Akinbami, L. J., Moorman, J. E., Garbe, P. L., & Sondik, E. J. (2009). Status of childhood asthma in the United States, 1980–2007. *Pediatrics*, 123 (Suppl. 3), 131-145. doi:10.1542/peds.2008-2233C
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., Text Revision). Washington DC: Author.
- Annesi-Maesano, I., Moreau, D., & Strachan, D. (2001). In utero and perinatal complications preceding asthma. *Allergy*, 56, 491-497. doi:10.1034/j.1398-9995.2001.056006491.x
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, 4, 561-571.
- Brown, G., Pennington, D., & Madrid, A. (1999). *Maternal-infant bonding survey*. Unpublished survey.
- Cassibba, R., van Ijzendoorn, M. H., Bruno, S., & Coppola, G. (2004). Attachment of mothers and children with recurrent asthmatic bronchitis. *Journal of Asthma*, 41(4), 419-431.
- Centers for Disease Control and Prevention. (2011, May). *Vital signs: Asthma in the U.S.* Retrieved from <http://www.cdc.gov/VitalSigns/pdf/2011-05-vitalsigns.pdf>
- DeFrances, C. J., Cullen, K. A., & Kozak, L. J. (2007). National hospital discharge survey: 2005 annual summary with detailed diagnosis and procedure data. *Vital Health Statistics* 13(165), 1-209. Abstract retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/18350768>
- Feinberg, S. (1988). *Degree of maternal infant bonding and its relationship to pediatric asthma and family environments* (Unpublished doctoral dissertation). The Professional School of Psychology, San Francisco, CA.
- Kero, J., Gissler, M., Gronlund, M. M., Kero, P., Koskinen, P., Hemminki, E., & Isolauri, E. (2002). Mode of delivery and asthma--Is there a connection? *Pediatric Research*, 52(1), 6-11.
- Klaus, M. H., & Kennell, J. H. (1976). *Maternal-infant bonding*. St. Louis, MO: C. V. Mosby.
- Klinnert, M. D., Nelson, H. S., Price, M. R., Adinoff, A. D., Leung, D. Y., & Mrazek, D. A. (2001, October). Onset and persistence of childhood asthma: Predictors from infancy. *Pediatrics*, 108(4), E69.
- Kozyrskij, A. L., Mai, X. M., McGrath, P., HayGlass, K. T., Becker, A. B., & MacNeil, B. (2008). Continued exposure to maternal distress in early life increases the risk of

- childhood asthma. *American Journal of Respiratory Critical Care Medicine*, 177, 142-147. doi:10.1164/rccm.200703-381OC
- Kozyrskyj, A. L., Mustard, C. A., & Becker, A. B. (2003). Childhood wheezing syndromes and healthcare data. *Pediatric Pulmonology*, 36, 131-136. doi:10.1002/ppul.10312
- Lange, N. E., Bunyavanich, S., Silberg, J. L., Canino, G., Rosner, B. A., & Celedon, J. C. (2011). Parental psychosocial stress and asthma morbidity in Puerto Rican twins. *Journal of Allergy and Clinical Immunology*, 127(3), 734-740.
- Lara, M., Rosenbaum, S., Rachelefsky, G., Nicholas, W., Morton, S. C., Emont, S., . . . Weiss, K. B. (2002). Improving childhood asthma outcomes in the United States: A blueprint for policy action. *Pediatrics*, 109(5), 919-930.
- Madrid, A. (2005/2006). Helping children with asthma by repairing maternal-infant bonding problems. *American Journal of Clinical Hypnosis*, 48(2,3), 199-211.
- Madrid, A., Ames, R., Horner, D., Brown, G., & Navarrette, L. (2004). Improving asthma symptoms in children by repairing the maternal-infant bond. *Journal of Prenatal and Perinatal Psychology and Health*, 18(3), 221-231.
- Madrid, A., Ames, R., Skolek, S., & Brown, G. (2000). Does maternal-infant bonding therapy improve breathing in asthmatic children? *Journal of Prenatal and Perinatal Psychology and Health*, 15(2), 90-112.
- Madrid, A., Skolek, S., & Shapiro, F. (2006). Repairing failures in bonding through EMDR. *Clinical Case Series*, 5(4), 271-286.
- Mantymaa, M., Puura, K., Luoma, I., Salmelin, R., Davis, H., Tsiantis, J., . . . Tamminen, T. (2003). Infant-mother interaction as a predictor of child's chronic health problems. *Child: Care, Health and Development*, 29(3), 181-191. doi:10.1046/j.1365-2214.2003.00330.x
- Martin, M. (2004). Asthma epidemic impacts Sonoma County. *Sonoma Medicine*, 55(1). Retrieved from <http://www.scma.org/magazine/?vol=55&num=1>
- Martinez, F. (2002). Development of wheezing disorders and asthma in preschool children. *Pediatrics*, 109(2), 362-367.
- Mead, V. (2004). A new model for understanding the role of environmental factors in the origins of chronic illness. *Medical Hypotheses*, 63(6), 1035-1046. doi:10.1016/j.mehy.2004.04.014
- Milam, J., McConnell, R., Yao, L., Berhane, K., Jerrett, M., & Richardson, J. (2008). Parental stress and childhood wheeze in a prospective cohort study. *Journal of Asthma*, 45, 319-323. doi:10.1080/02770900801930277
- Mrazek, D. A., Klinnert, M. D., Mrazek, P., & Macey, T. (1991). Early asthma onset: Consideration of parenting issues. *Journal of the American Academy of Child & Adolescent Psychiatry*, 30(2), 277-282. doi:10.1097/00004583-199103000-00017
- Nafstad, P., Samuelsen, S. O., Irgens, L. M., & Bjerkedal, T. (2003). Pregnancy complications and the risk of asthma among Norwegians born between 1967 and 1993. *European Journal of Epidemiology*, 18, 755-761. doi:10.1023/A:1025395405101
- Pennington, D. (1991). *Events associated with maternal-infant bonding deficits and severity of pediatric asthma* (Unpublished doctoral dissertation). The Professional School of Psychology, San Francisco, CA.
- Roduit, C., Scholtens, S., de Jongste, J. C., Gerritsen, J., Postma, D. S., Brunekreef, B., . . . Smit, H. A. (2009). Asthma at 8 years of age in children born by caesarean section. *Thorax*, 64(2), 107-113. doi:10.1136/thx.2008.100875

- Schwartz, M. (1988). *Incidence of events associated with maternal-infant bonding disturbances in a pediatric asthma population* (Unpublished doctoral dissertation), Rosebridge Graduate School, Walnut Creek, CA.
- U.S. Department of Health and Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute. (2003). *NAEPP expert report: Guidelines for the diagnosis and management of asthma* (NIH Publication No. 02-5074). Retrieved from http://msdh.ms.gov/msdhsite/_static/resources/2100.pdf
- Wright, R. J., Cohen, S., Carey, V., Weiss, S. T., & Gold, D. R. (2002). Parental stress as a predictor of wheezing in infancy: A prospective birth-cohort study. *American Journal of Respiratory Critical Care Medicine*, *165*(3), 358-365.
- Wright, R. J., Visness, C. H., Calatroni, A., Grayson, M. H., Gold, D. R., Sandel, M. T., . . . Gern, J. E. (2010). Prenatal maternal stress and cord blood innate and adaptive cytokine responses in an inner-city cohort. *American Journal of Respiratory Critical Care Medicine*, *182*(1), 25-33. doi:10.1164/rccm.200904-0637OC

APPENDIX A

MATERNAL-INFANT BONDING SURVEY

Name _____ Child _____

The following questions are to see if there are any of factors that may have had some impact on your child's birth.

POSSIBLE PREGNANCY PROBLEMS Y N ?

Had worrisome bleeding during pregnancy

Had toxemia

Vomited a lot

Had to be medicated

Gained too much weight

Took a lot of illegal drugs

Drank excessively

Was sick through much of pregnancy

Labor lasted longer than 15 hours

Had a difficult delivery

Had a Caesarean Section

Was put to sleep for delivery

Got hurt during pregnancy

POSSIBLE OTHER PREGNANCY PROBLEMS Y N ?

Had a previous miscarriage

Was overly depressed during pregnancy

Was very scared during pregnancy

Lost someone close during pregnancy

Had marital problems during pregnancy

Had serious financial problems during pregnancy

Had a serious loss after the child was born

Was overly depressed after the child was born

Had emotional problems after the child was born

Was very sick during delivery

Was very sick after the baby was born

Child was a twin or triplet

Moved during pregnancy or first year

HOW WAS YOUR RELATIONSHIP TO THE BABY'S FATHER DURING PREGNANCY?

WHAT HAPPENED TO THE BABY AFTER IT WAS BORN?

HOW LONG AFTER THE BABY WAS BORN DID YOU HOLD IT?

WHAT WAS IT LIKE WHEN YOU FIRST HELD THE BABY?

BABY'S CONDITION	Y	N	?
------------------	---	---	---

- | | | | |
|---|--|--|--|
| Injured during birth | | | |
| Was born jaundiced | | | |
| Had trouble breathing | | | |
| Born with cord around neck | | | |
| Was sick after birth | | | |
| Spent time in an incubator | | | |
| Spent time in an Intensive Care Nursery | | | |
| Was born premature | | | |
| Had an infection | | | |
| Needed oxygen | | | |
| Vomited often | | | |
| Gagged often | | | |
| Was kept in hospital after mother went home | | | |
| Was born breech | | | |

LIST OTHER PROBLEMS THAT OCCURRED DURING PREGNANCY, DELIVERY, OR THE FIRST COUPLE OF MONTHS OF THE BABY'S LIFE.

- 1. _____
- 2. _____
- 3. _____

BABY'S PROBLEMS IN THE BEGINNING Y N ?

- Colic
- Coughed a lot
- Wheezed
- Was sickly
- Difficult to calm or comfort
- Cried often
- Was demanding
- Could not be alone
- Did not like to be held
- Irritable
- Was easily upset
- Had lots of mucous
- Was frightened easily
- Seemed in pain a lot
- Difficult to console
- Feeding difficulties
- Was not affectionate

ANY OTHER THOUGHTS ABOUT YOUR BABY IN THE FIRST YEAR OF LIFE?

ANY OTHER THOUGHTS ABOUT YOU OR YOUR FAMILY DURING PREGNANCY, BIRTH, OR THE FIRST YEARS OF YOUR CHILD'S LIFE?

ANY OTHER THOUGHTS ABOUT YOUR CHILD'S CONDITION?

Thank you!

APPENDIX B

CHILD INFORMATION

CHILD'S NAME _____ AGE _____

Parent's name _____

This questionnaire asks lots of questions about your child's condition. Please answer the best you can, even though you may contradict yourself in places. It's all right. We'll get the idea of your child's health through all of this.

Age when child was diagnosed with asthma _____

Age when symptoms began: _____

Who made the diagnosis? _____

How did this occur? _____

Course of asthma: _____

What are your child's triggers? _____

Child's symptoms? _____

What medications is your child taking, how much and how often?

Your child's daytime symptoms are:

- None 2 days a week or less
 greater than twice a week
 daily continual

Your child's night time symptoms are:

- none 2 nights or less a month
 more than 2 nights a month
 more than once a week frequent

How many doctor visits for asthma in the last 6 months? ____

How many Emergency Room visits in the last 6 months? ____

How many hospitalizations in the last 6 months? ____ Total days? ____

How many times did your child use oral steroids in the last 6 months? ____ Total days ____

Days absent from school or housebound last 6 months? ____

Wheezes with exercise? never sometimes frequently
 almost always always ____

Child's overall health (A-F): ____

Child's overall energy level (A-F): ____

Please check ONE answer for each the the questions.

1. In the past 4 weeks, how much of the time did your child's asthma keep him/her from getting much done at school or at home?

- none of the time
 a little of the time
 some of the time
 most of the time
 all of the time

2. During the past 4 weeks, how often has your child had shortness of breath?

- not at all
- once or twice a week
- 3-6 times a week
- once a day
- more than once a day

3. During the past 4 weeks, how often did your child's asthma symptoms (wheezing, coughing, shortness of breath, chest tightness or pain) wake you up at night or earlier than usual in the morning?

- not at all
- once or twice
- once a week
- 2 to 3 times a week
- 4 or more nights a week

4. During the past 4 weeks, how often has your child used the rescue inhaler or nebulizer medication (such as albuterol)?

- not at all
- once a week or less
- a few times a week
- 1 or 2 times a week
- 3 or more times a week

5. How would you rate your child's asthma control during the past 4 weeks?

- completely controlled
- well controlled
- somewhat controlled
- poorly controlled
- not controlled at all

Thank you!